



TANBead® Nucleic Acid Extraction Kit

Plasmid Extraction Auto Plate

(For use with the Maelstrom 9600 series)

RUO

W6PEA46

(For Professional Use Only) V1

1. Intended Use

This product is designed for isolating nucleic acid from various samples, which can be performed by using TANBead® Nucleic Acid Extractor and is intended for research use only.

2. Purpose

TANBead® Nucleic Acid Extraction Kit (W6PEA46) is designed for a simple and convenient method of isolating high-quality plasmid DNA from *E. coli* strains including DH10B, DH5α, BL21(DE3), and TOP10. This kit, with Maelstrom 9600 series, utilizes the unique non-alkaline lysis method to simplify the pre-treatment protocol of plasmid extraction. It has no repetitive centrifugation steps, reducing time for manual processing and error rate. The purified plasmid can be directly analyzed (such as Nanodrop, PCR, agarose gel electrophoresis, etc.) and used for downstream experiments.

3. The basic principle

The silicon dioxide layer coated on the magnetic beads can adsorb the negatively charged molecules to purify nucleic acids from samples.

4. Specification

Starting Materials	<i>E. coli</i> culture (OD ₆₀₀ 1~5)
Elution Volume	50 - 80 µL
Typical DNA yield	Up to 10 µg
Typical A260/A280	1.7 - 2.0

5. Component Supplied with the Kit



Auto Plate	5	Auto Plate with reagent buffers
Incubation Buffer	35 mL x 1	Tris buffer, surfactants
Elution Buffer	1.5 mL x 1	Nuclease-Free Water
Spin tips	96 tips	Spin tip assembled box
RNase A	100 µL x 1	-
Lysozyme	1	Add 100 µL sterile water before use
Protocol	1	Instruction manual

6. Auto Plate Content and Plate Position

Plate Position	Buffer	Volume (µL)
1	Binding Buffer	600
2	Washing Buffer 1	800
3	Washing Buffer 2	800
4	-	-
5	Magnetic Beads	800
6	Elution Buffer	80
7	-	-
8	Spin tip	-

7. Kit Storage and Shelf Life

- Components under room temperature (15 ~ 35°C) can be stored until the expiration date labeled on the box.
- The RNase A and Lysozyme are transported at room temperature. Upon received, please store RNase A and Lysozyme at 2~8°C.
- Incubation buffer should be stored in 4°C after the addition of Lysozyme and RNase A.

8. Precautions

- For research use only.
- Avoid using expired reagents.
- When the temperature is below 20°C, place the reagent Plate in an oven (preheated 42 ~ 60°C) from 5 to 10 minutes.
- Avoid vigorous shaking, in order to avoid excessive formation of foam.
- Carefully remove aluminum foil to avoid splashing.
- Do not expose the opened reagents or Auto Plates to air. The evaporation would lead to pH change, or effect on the extraction effectiveness.

- Please check the integrity of the Auto Plates and remember to mount the spin tips into the appropriate position of the suitable instrument before operating them.
- Please wear a mask and disposable gloves when handling.
- Use sterile consumables to avoid nuclease contamination.
- Reagent solution contains guanidine salt, avoid using bleach containing detergent.
- Avoid eyes, skin, and clothing contact with reagents. In case of any contact, flush with flowing water.
- If any serious incident occurs, please report to the manufacturer and the competent authority of the member state in which the user and/or the patient is established.

9. Materials required, Not Supplied

- TANBead® Nucleic Acid Extraction System
Model: Maelstrom 9600 series (non-sterile)
- Disposable gloves
- Micropipette, disposable tips (10 µL/ 200 µL/ 1000 µL)
- 1.5 mL microcentrifuge tube
- 15 mL / 50 mL conical tube

10. Sample Collection, Transportation, and Storage

■ Sample collection

- The plasmid recovery is directly affected by host strain, antibiotic selection, culture broth, plasmid copy number, plasmid size, insert toxicity, etc. Handling the transformation and enrichment of *E. coli* properly to ensure maximum plasmid yield.
- We suggest using LB (Luria Bertani) as the culture medium for plasmid isolation. The best starting biomass of *E. coli* is an OD₆₀₀ of 1~5, which should be incubated overnight at 37°C.
- We do not recommend using *E. coli* culture during cold storage as material, it may lead to the contamination of genomic DNA.

11. Nucleic Acids Extraction Protocol

- Add all Lysozyme and RNase A into Incubation Buffer.
- Harvest the overnight culture of *E. coli* into a 1.5 mL tube by centrifugation at 6,000 × *g* for 2 min, then carefully remove the supernatant.
- Add 300 µL incubation buffer into a 1.5 mL tube and resuspended the pellet.
- Incubate the sample at 65°C for 5 min.
- Centrifuged at 13,000 × *g* for 3 min.
- Carefully remove the aluminum foil on the Auto Plates.
- Transfer 300 µL suspension sample into **plate #1** of Auto Plate (Plate filled with binding buffer).
- Select the program "**6PE**". The parameters are given in the following section.
- Follow the guide shown on the screen and place plates carefully. Make sure that the chamfer of the plate faces toward the same direction.
- Carefully remove the Auto Plates when the program is finished.
- Use micropipette to transfer the purified nucleic acids from **plate #6** to a clean tube.
- Discard used Auto Plates and spin tips into the waste recycling bin.

12. Program

■ Maelstrom 9600

Program Name: 6PE								
Plate	1	2	3	4	5	6	7	8
Volume(μL)	900	800	800	-	800	100	-	-
Keep Temp.	60	0	0	-	-	0	-	-
Action	For. U/D	For.	For.	For.	For.	For.	-	-
Name	LB	WB1	WB2	WB2	MB	EB	-	TIP
Step	Plate	Temp. (°C)	Mixing (min)	Mixing (rpm)	Collect (sec)	Vapor (min)	Pause	
1	5	-	0	2500	12	0	Off	
2	1	60	3	2500	12	0	Off	
3	2	0	1	3000	12	0	Off	
4	3	0	1	3000	12	0	Off	
5	5	-	1	3000	12	5	Off	
6	6	60	5	2000	60	0	Off	

⚠ Temperature set as "0" represents room temperature!

■ Maelstrom 9610

Program Name: 6PE								
Plate	1	2	3	4	5	6	7	8
Volume(μL)	900	800	800	-	800	100	-	-
Preheat.	60	25	25	-	-	25	-	-
Action	For. U/D	For.	For.	For.	For.	For.	-	-
Name	LB	WB1	WB2	WB2	MB	EB	-	TIP
Step	Plate	Temp. (°C)	Mixing (min)	Mixing (rpm)	Collect (sec)	Vapor (min)	Pause	
1	5	-	0	2500	12	0	Off	
2	1	60	3	2500	12	0	Off	
3	2	25	1	3000	12	0	Off	
4	3	25	1	3000	12	0	Off	
5	5	-	1	3000	12	5	Off	
6	6	60	5	2000	60	0	Off	

⚠ Temperature set as "25" represents room temperature!

13. Result

Total nucleic acid yield and purity were detected using Nanodrop spectrophotometers: Up to 10 μg DNA per test and the A260/A280 ratio of the nucleic acid is between 1.7 to 2.0.

14. Reagent performance

■ Host strain

Several *E. coli* strains including DH10B, DH5α, BL21(DE3), and TOP10 can be used for plasmid isolation by this kit.

■ Plasmid type

Common cloning vector plasmids (such as pUC, pBS, pBR and pET) can be isolated by this kit.

■ Plasmid size

The plasmid which is less than < 15 kb can be isolated by this kit, isolating larger size of plasmid may decrease the yield of plasmid.

■ Extraction time and throughput

Less than 28 min / 96 preps on Maelstrom 9600 series

15. Explanation of Symbols

	Manufacturer		Consult instructions for use
	Temperature limit		Contains sufficient for test
	Catalogue number		Caution
	Batch code		Non-sterile
	Do not re-use		Keep away from sunlight
	Date of manufacture		Use-by date
	For research use only		

